

DAVID L. HOOVER, M.D.*; AND ARTHUR M. FRIEDLANDER, M.D.†

*C₁, C₂, A₁; D₁, B₁, D₁, A₁, I₁, D. C. 20307-5100; A₁, H₁, B₁, 20814-4799
†C₁, C₂, A₁; C₂, B₁, D₁, A₁, I₁, I₁, D₁, F₁, D₁, E₁, 21702-5011; C₁, A₁, B₁, H₁, 4301

suffering from contagious abortion.⁶ In 1917, A. C. Evans recognized that Bang's organism was identical to that described by Bruce as the causative agent of human brucellosis. The organism infects mainly cattle, sheep, goats, and other ruminants, in which it causes abortion, fetal death, and genital infection.^{7,8} Humans, who are usually infected incidentally by contact with infected animals or ingestion of dairy foods, may develop numerous symptoms in addition to the usual ones of fever, malaise, and muscle pain. Disease frequently becomes chronic and may relapse, even with treatment.

The ease of transmission by aerosol suggests that *B. melitensis* organisms might be a candidate for use as a biological warfare agent. Indeed, the United States began development of *B. melitensis* as a biological weapon in 1942. The agent was formulated to maintain long-term viability, placed into bombs, and tested in field trials during 1944–1945 using animal targets. By 1967, the United States terminated its offensive program for development and deployment of *B. melitensis* as a biological weapon. Although the munitions developed were never used in combat, the studies reinforced the concern that *B. melitensis* organisms might be used against U.S. troops as a biological warfare agent.⁹

Brucellae are small, nonmotile, nonsporulating, nontoxigenic, nonfermenting, aerobic, Gram-negative coccobacilli that may, based on DNA homology, represent a single species.¹⁰ Conventionally, however, they are classified into six species, each comprising several biovars. Each species has a characteristic, but not an absolute, predilection to infect certain animal species (Table 25-1). Only *B. melitensis*, *B. abortus*, *B. canis*, and *B. maritima* cause disease in man. Infection of humans with *B. melitensis* and *B. abortus* has not been described.

Brucellae grow best on trypticase, soy-based, or other enriched media with a typical doubling time of 2 hours. Most biovars of *B. melitensis* are

thionine or basic fuchsin dyes; agglutination by antisera directed against certain lipopolysaccharide epitopes; and by susceptibility to lysis by bacte-

riophage. Recently, analysis of fragment lengths of deoxyribonucleic acid (DNA) cut by various restriction enzymes has also been used to differentiate brucellae groupings.¹⁰

The lipopolysaccharide (LPS) component of the outer cell membranes of brucellae is quite different—both structurally and functionally—from that of other Gram-negative organisms.^{11,12} The lipid A portion of a B \square organism LPS contains fatty acids 16 carbons long, and lacks the 14-carbon myristic acid typical of lipid A of Enterobacteriaceae. This unique structural feature may underlie

the remarkably reduced pyrogenicity (less than 1/100th) of B \square LPS, compared with the pyrogenicity of *E. coli* LPS.¹³ In addition, the O-polysaccharide portion of LPS from smooth organisms contains an unusual sugar, 4,6-dideoxy-4-formamido- α -D-mannopyranoside, which is expressed either as a homopolymer of α -1,2-linked sugars (A type), or as 3 α -1,2 and 2 α -1,3-linked sugars (M type). These variations in O-polysaccharide linkages lead to specific, taxonomically useful differences in immunoreactivity between A and M sugar types.¹⁴

Animals may transmit B \square organisms during septic abortion, at the time of slaughter, and in their milk. Brucellosis is rarely, if ever, transmitted from person to person. The incidence of human disease is thus closely tied to the prevalence of infection in sheep, goats, and cattle, and to practices that allow exposure of humans to potentially infected animals or their products. In the United States, where most states are free of infected animals and where dairy products are routinely pasteurized, illness occurs primarily in individuals such as veterinarians, shepherds, cattlemen, and slaughterhouse workers who have occupational exposure to infected animals. In many other countries, humans more commonly acquire infection by ingestion of unpasteurized dairy products, especially cheese.

Less obvious exposures can also lead to infection. In Kuwait, for example, disease with a relatively

A C B □ □

in lymph nodes, liver, spleen, mammary gland, joints, kidneys, and bone marrow.

In macrophages, brucellae may inhibit fusion of phagosomes and lysosomes, and replicate in the phagosome.²³ If unchecked by macrophage microbicidal mechanisms, the bacteria destroy their host cells and infect additional cells. Brucellae can also replicate extracellularly in host tissues. Histopathologically, the host cellular response may range from abscess formation to lymphocytic infiltration to granuloma formation with caseous necrosis.

Studies in experimental models have provided important insights into host defenses that eventually control infection with *B*□ organisms. Se-

A C B □ □

ganism does not have O-polysaccharide on its surface. Immunoenzymatic assays (eg, enzyme-linked immunosorbent assays [ELISAs]) have been developed for use with *B. melitensis*, but are not well standardized. ELISAs developed for other brucellae similarly suffer from lack of standardization.

In addition to serologic testing, diagnosis should

axial skeleton are favored targets; arthritis appears in approximately one third of patients. Fatalities occur rarely, usually in association with central nervous system or endocardial infection.

Serologic diagnosis uses an agglutination test that detects antibodies to lipopolysaccharide. This test, however, is not useful to diagnose infection caused by *B. abortus*, a naturally O-polysaccharide deficient strain. Infection can be most reliably con-

firmed by culture of blood, bone marrow, or other infected body fluids, but the sensitivity of culture varies widely.

Nearly all patients respond to a 6-week course of oral therapy with a combination of rifampin and doxycycline; fewer than 10% of patients relapse. Six weeks of doxycycline with addition of streptomycin for the first 3 weeks is also effective therapy. No vaccine is available for humans.

REFERENCES

1. Evans AC. Comments on the early history of human brucellosis. In: Larson CH, Soule MH, eds. *Brucellosis*. Baltimore, Md: Waverly Press; 1950: 1-8.
2. Cleghorn G. *History of the Brucellae* (Edinburgh 1744-1749). London, England; 1751. Cited in: Evans AC. Comments on the early history of human brucellosis. In: Larson CH, Soule MH, eds. *Brucellosis*. Baltimore, Md: Waverly Press; 1950: 1-8.
3. Marston JA. Report on fever (Malta). *Amer J Hyg*. 1861;3:486-521. Cited in: Evans AC. Comments on the early history of human brucellosis. In: Larson CH, Soule MH, eds. *Brucellosis*. Baltimore, Md: Waverly Press; 1950: 1-8.
4. Bruce D. Note on the discovery of a micro-organism in Malta fever. *Lancet* (London). 1887;39:161-170. Cited in: Evans AC. Comments on the early history of human brucellosis. In: Larson CH, Soule MH, eds. *Brucellosis*. Baltimore, Md: Waverly Press; 1950: 1-8.
5. Hughes ML. *Brucella*. London, England: Macmillan and Co; 1897. Cited in: Evans AC. Comments on the early history of human brucellosis. In: Larson CH, Soule MH, eds. *Brucellosis*. Baltimore, Md: Waverly Press; 1950: 1-8.
6. Bang B. Die Aetiologie des seuchenhaften ("infectiösen") Verwerfens. *Monatsschr Med Naturg* (Jena). 1897;1:241-278. Cited in: Evans AC. Comments on the early history of human brucellosis. In: Larson CH, Soule MH, eds. *Brucellosis*. Baltimore, Md: Waverly Press; 1950: 1-8.
7. Meador VP, Hagemoser WA, Deyoe BL. Histopathologic findings in *Brucella abortus*-infected, pregnant goats. *Am J Vet Res*. 1988;49(2):274-280.
8. Nicoletti P. The epidemiology of bovine brucellosis. *Am J Hyg*. 1980;24(69):69-98.
9. Department of the Army. *Antigenic Properties of Brucella abortus*, Vols 1 and 2. Washington, DC: HQ, DA; 24 February 1977. Unclassified.
10. Grimont F, Verger JM, Cornelis P, et al. Molecular typing of *Brucella abortus* with cloned DNA probes. *J Clin Microbiol*. 1992;143(1):55-65.
11. Bundle DR, Cherwonogrodzky JW, Caroff M, Perry MB. The lipopolysaccharides of *Brucella abortus* and *Brucella melitensis*. *J Biol Chem*. 1987;138(1):92-98.
12. Moreno E, Borowiak D, Mayer H. *Brucella abortus* lipopolysaccharides and polysaccharides. *J Biol Chem*. 1987;138(1):102-105.
13. Goldstein J, Hoffman T, Frasch C, et al. Lipopolysaccharide (LPS) from *Brucella abortus* is less toxic than that from *Escherichia coli*, suggesting the possible use of *Brucella abortus* or LPS from *Brucella abortus* as a carrier in vaccines. *J Biol Chem*. 1992;60(4):1385-1389.
14. Cherwonogrodzky JW, Perry MB, Bundle DR. Identification of the A and M antigens of *Brucella abortus* as the O-polysaccharides of smooth lipopolysaccharides. *J Biol Chem*. 1987;33(11):979-981.

15. Mousa AR, Elhag KM, Khogali M, Marafie AA. The nature of human brucellosis in Kuwait: Study of 379 cases. *I, D* . 1988;10(1):211-217.
16. Robson JM, Harrison MW, Wood RN, Tilse MH, McKay AB, Brodribb TR. Brucellosis: Re-emergence and changing epidemiology in Queensland. *J A* . 1993;159(3):153-158.
17. Dajani YF, Masoud AA, Barakat HF. Epidemiology and diagnosis of human brucellosis in Jordan. *J □ H* . 1989;92(3):209-214.
18. Mousa AM, Elhag KM, Khogali M, Sugathan TN. Brucellosis in Kuwait: A clinico-epidemiological study. □, □ *H* . 1987;81(6):1020-1021.
19. Buchanan TM, Hendricks SL, Patton CM, Feldman RA. Brucellosis in the United States, 1960-1972: An abattoir-associated disease, III: Epidemiology and evidence for acquired immunity. *(B □ □)*. 1974;53(6):427-439.
20. Ackermann MR, Cheville NF, Deyoe BL. Bovine ileal dome lymphoepithelial cells: Endocytosis and transport of B□ □ strain 19. *□* . 1988;25(1):28-35.
21. Elsbach P. Degradation of microorganisms by phagocytic cells. *I, D* . 1980;2:106-128.
22. Braude AI. Studies in the pathology and pathogenesis of experimental brucellosis, II: The formation of the hepatic granulomas and its evolution. *J I, D* . 1951;89:87-94.
23. Harmon BG, Adams LG, Frey M. Survival of rough and smooth strains of B□ □ in bovine mammary gland macrophages. *A J* . 1988;49(7):1092-1097.
24. Young EJ, Borchert M, Kretzer FL, Musher DM. Phagocytosis and killing of B□ □ by human polymorphonuclear leukocytes. *J I, D* . 1985;151(4):682-690.
25. Corbeil LB, Blau K, Inzana TJ, et al. Killing of B□ □ by bovine serum. *I, L □* . 1988;56(12):3251-3261.
26. Montaraz JA, Winter AJ, Hunter DM, Sowa BA, Wu AM, Adams LG. Protection against B□ □ in mice with O-polysaccharide-specific monoclonal antibodies. *I, L □* . 1986;51(3):961-963.
27. Araya LN, Elzer PH, Rowe GE, Enright FM, Winter AJ. Temporal development of protective cell-mediated and humoral immunity in BALB/c mice infected with B□ □ . *J L □* . 1989;143(10):3330-3337.
28. Zhan Y, Cheers C. Endogenous gamma interferon mediates resistance to B□ □ infection. *I, L □* . 1993;61(11):4899-4901.
29. Jiang X, Baldwin CL. Effects of cytokines on intracellular growth of B□ □ . *I, L □* . 1993;61(1):124-134.
30. Anderson TD, Cheville NF. Ultrastructural morphometric analysis of B□ □ -infected trophoblasts in experimental placentitis: Bacterial replication occurs in rough endoplasmic reticulum. *A J □* . 1986;124(2):226-237.
31. Anderson TD, Cheville NF, Meador VP. Pathogenesis of placentitis in the goat inoculated with B□ □ , II: Ultrastructural studies. *□* . 1986;23(3):227-239.
32. Young EJ. Human brucellosis. *I, D* . 1983;5(5):821-842.
33. Crosby E, Llosa L, Miro QM, Carrillo C, Gotuzzo E. Hematologic changes in brucellosis. *J I, D* . 1984;150(3):419-424.

34. Gotuzzo E, Alarcon GS, Bocanegra TS, et al. Articular involvement in human brucellosis: A retrospective analysis of 304 cases. *J Infect Dis*. 1982;12(2):245–255.
35. Alarcon GS, Bocanegra TS, Gotuzzo E, Espinoza LR. The arthritis of brucellosis: A perspective one hundred years after Bruce's discovery. *J Infect Dis*. 1987;14(6):1083–1085.
36. Mousa AR, Muhtaseb SA, Almudallal DS, Khodeir SM, Marafie AA. Osteoarticular complications of brucellosis: A study of 169 cases. *J Infect Dis*. 1987;9(3):531–543.
37. Howard CB, Alkrinawi S, Gadalia A, Mozes M. Bone infection resembling phalangeal microgeodic syndrome in children: A case report. *J Hyg (Lond)*. 1993;18(4):491–493.
38. Rotes-Querol J. Osteo-articular sites of brucellosis. *Am J Trop Med Hyg*. 1957;16:63–68.
39. Ibrahim AIA, Shetty SD, Saad M, Bilal NE. Genito-urinary complications of brucellosis. *Bull WHO*. 1988;61:294–298.
40. Kelalis PP, Greene LF, Weed LA. Brucellosis of the urogenital tract: A mimic of tuberculosis. *JAMA*. 1962;88:347–353.
41. Lubani MM, Dudin KI, Sharda DC, et al. Neonatal brucellosis. *Eur J Clin Invest*. 1988;14(5):520–522.
42. Buchanan TM, Faber LC, Feldman RA. Brucellosis in the United States, 1960–1972: An abattoir-associated disease, I: Clinical features and therapy. *Am J Hyg (B)*. 1974;53(6):403–413.
43. Peery TM, Belter LF. Brucellosis and heart disease, II: Fatal brucellosis. *Am J Hyg*. 1960;36:673–697.
44. Young EJ. Serologic diagnosis of human brucellosis: Analysis of 214 cases by agglutination tests and review of the literature. *J Infect Dis*. 1991;13(3):359–372.
45. Buchanan TM, Faber LC. 2-mercaptoethanol *B* agglutination test: Usefulness for predicting recovery from brucellosis. *J Clin Microbiol*. 1980;11(6):691–693.
46. Gotuzzo E, Carrillo C, Guerra J, Llosa L. An evaluation of diagnostic methods for brucellosis—The value of bone marrow culture. *J Infect Dis*. 1986;153(1):122–125.
47. Hall WH. Modern chemotherapy for brucellosis in humans. *J Infect Dis*. 1990;12(6):1060–1099.
48. Luzzi GA, Brindle R, Sockett PN, Solera J, Klenerman P, Warrell DA. Brucellosis: Imported and laboratory-acquired cases, and an overview of treatment trials. *J Infect Dis*. 1993;87(2):138–141.
49. Joint FAO/WHO expert committee on brucellosis. ~~Report of the expert committee on brucellosis~~. *Wkly Rep WHO*. 1986;740(1):1–132.
50. Ariza J, Gudiol F, Pallares R, et al. Treatment of human brucellosis with doxycycline plus rifampin or doxycycline plus streptomycin: A randomized, double-blind study. *Am J Trop Med Hyg*. 1992;117(1):25–30.
51. Montejo JM, Alberola I, Glez ZP, et al. Open, randomized therapeutic trial of six antimicrobial regimens in the treatment of human brucellosis. *Clin Infect Dis*. 1993;16(5):671–676.
52. Chan R, Hardiman RP. Endocarditis caused by *B*. *JAMA*. 1993;158(9):631–632.
53. Lubani MM, Dudin KI, Sharda DC, et al. A multicenter therapeutic study of 1,100 children with brucellosis. *J Infect Dis*. 1989;8(2):75–78.